

# Modeling reveals the role of coastal upwelling and hydrologic inputs on biologically distinct water exchanges in a Great Lakes estuary

Qianqian Liu<sup>a,b</sup>, Eric J. Anderson<sup>c,\*</sup>, Yinglong Zhang<sup>d</sup>, Anthony D. Weinke<sup>a</sup>, Katie L. Knapp<sup>a</sup>, Bopaiah A. Biddanda<sup>a</sup>

<sup>a</sup> Annis Water Resources Institute, Grand Valley State University, Muskegon, MI, 49441, USA

<sup>b</sup> Cooperative Institute for Great Lakes Research, University of Michigan, Ann Arbor, MI, 48108, USA

<sup>c</sup> Great Lakes Environmental Research Laboratory, National Oceanic and Atmospheric Administration, Ann Arbor, MI, 48108, USA

<sup>d</sup> Virginia Institute of Marine Science, Gloucester Point, VA, 23062, USA

## ARTICLE INFO

### Keywords:

Freshwater estuary  
Cold-water intrusion  
Upwelling  
Hydrologic shift  
Numerical model  
Hypoxia

## ABSTRACT

Freshwater estuaries everywhere are under stress from anthropogenic activities and climate change. Muskegon Lake Estuary (MLE) is a freshwater estuary along the eastern shore of Lake Michigan characterized by algal blooms and hypoxia during the summer and designated as an Area of Concern (AOC) by the EPA. We developed a 3-D hydrodynamic model using the Semi-implicit Cross-scale Hydrosystems Integrated System Model (SCHISM) to study the hydrodynamics of MLE with a focus on the cold-water intrusions from Lake Michigan into MLE. Substantial water exchange process was validated by comparisons with observations in the near-shore region of Lake Michigan and in the navigation channel between Lake Michigan and MLE. The model found that the cold-water intrusions from Lake Michigan to MLE occur during summer stratification, amounting to as much as 10% of MLE's total volume during one single episodic event. The intrusion was accompanied by a stronger surface outflow in the opposite direction, which may accelerate the delivery of MLE water to Lake Michigan. Through process-oriented model experiments, we examined the cold-water intrusion's responses to hydrological shift under climate change, and found that the increase in riverine input during upwelling weakens the intrusion. In addition, an increase of navigation channel width strengthens the cold-water intrusion, and that intrusion strength as well as intrusion period was directly related to wind speed. Our observation-modeling based findings would provide a good reference for the future study of biophysical interactions between coastal ocean and estuaries.

## 1. Introduction

Estuaries are an ecologically active interface between riverine systems and coastal waters, which provide unique habitats for many varieties of plants and animals, are highly productive systems, and undergo several complex biological and physical processes. Similar to brackish estuarine systems, freshwater estuaries involve mixing of chemically and biologically distinct waters, as they receive hydrologic inputs with carbon and nutrients from terrestrial sources and heavily urbanized population centers, and as a result can have large spatial variability in dissolved oxygen, conductivity, and turbidity (Fisher et al., 2015). Such land-water interface ecosystems are greatly affected by anthropogenic activities and climate change (Borja et al., 2010).

Though extremely large, the Laurentian Great Lakes, are quite vulnerable to anthropogenic stressors and climate driven changes in physical or biogeochemical conditions such as thermal stratification,

precipitation and runoff, and timing of the hydrometeorological cycle (Allan et al., 2012; Larson et al., 2013; Cotner et al., 2017). In addition, lakes are highly reactive sites for carbon metabolism and play a critically important role in the global carbon cycle because of the enhanced rates of carbon respiration and burial within them (Cole et al., 2007; Tranvik et al., 2009; Biddanda, 2017). As climate projections reveal intensification of extreme precipitation events over the next century (Donat et al., 2016; Wang et al., 2017), and subsequent shifts in riverine inputs to estuarine systems, an important concern arises regarding the impact of these changes on estuarine function – such as rising coastal eutrophication worldwide (Sinha et al., 2017).

Extensive studies have focused on the mechanisms that control estuarine dynamics, including the exchange flow in different brackish estuarine systems (Chen et al., 2012), the generation of estuarine front (Geyer and Ralston, 2015; Liu et al., 2016a; b), turbulent mixing in estuaries and the resulting variations in water exchange (Simpson et al.,

\* Corresponding author.

E-mail address: [eric.j.anderson@noaa.gov](mailto:eric.j.anderson@noaa.gov) (E.J. Anderson).

1990; Rong and Li, 2012; Liu et al., 2017), and biologically distinct water exchange during coastal upwelling (Roegner et al., 2011). Roegner et al. (2011) found that the coastal upwelling supplies oxygen-depleted water to the Columbia River estuary and therefore deteriorates the estuarine habitats. Similarly, upwelling along the central California coast supports Harmful Algal Blooms (HABs) in some open embayments by delivering nutrients from the bottom of the coastal ocean to the embayments but sustaining its stratification (Pitcher et al., 2010). Along the California coast, in 2007, an inshore anoxia over a 5 km stretch of coastline at Erendira is believed to be caused by a strong upwelling region off Baja California, killed many animals, including tons of lobsters (Levin et al., 2009). Intrusions of saline hypoxic water to estuaries are commonly found, for example, from the ocean to Puget Bay (Deppe et al., 2017), from Gulf of Mexico to its embayments (Rabalais et al., 2002), and from Chesapeake Bay to the lower reaches of the Choptank River (Breitburg, 1992, 2002; Sanford et al., 1990). Similar intrusions of cold, dense oxygenated waters have also been found to occur in the freshwater estuaries of the Great Lakes; e.g., Lake Michigan water into Green Bay (Grunert, 2013) and Lake Ontario water into Hamilton Harbor (Lawrence et al., 2004), and the Muskegon Lake Estuary (MLE; Biddanda et al., 2018), the investigation area in this study.

In the Great Lakes, where tides are weak, winds become the dominant mechanism for water exchange between the Great Lakes and freshwater estuaries. However, there is little information on the drivers of exchange in a freshwater estuary and in particular, how the mechanisms that control important ecological variables such as dissolved oxygen are impacted by shifts in precipitation or meteorology due to a changing climate. To investigate the dominant mechanisms behind freshwater estuary exchange and the sensitivity of spatiotemporal variability of dissolved oxygen to these controls, this study focuses on the MLE, one of the several drowned river-mouth estuaries along the eastern coastline of Lake Michigan (Larson et al., 2013).

## 2. Study area

The MLE is a mesotrophic drowned river mouth along the eastern shore of Lake Michigan (Fig. S1 in Supplementary Material), with a surface area of 17 km<sup>2</sup>, a water volume of 119 million m<sup>3</sup>, a mean water depth of 7 m, a maximum depth of 21 m (Fig. 1; Steinman et al., 2008; Marko et al., 2013) and an average hydraulic residence time of 23 days. The estuary drains the second largest watershed in Michigan (approximately 7302 km<sup>2</sup>) including 53.2% forested, 23.0% agricultural, and 4.2% urban lands (Marko et al., 2013), and connects to an oligotrophic Lake Michigan through a 1 km long and 100 m wide navigation channel.

Historical observations show that the MLE is under the effects of anthropogenic activities and climate change (Steinman et al., 2008). The proportion of urbanized land within the Muskegon River watershed is projected to increase by 11.5% by the year 2040 (Tang et al., 2005), which changes the nature and quantity of nutrients, major ions, and dissolved organic carbon (DOC) inputs to the estuary. The US Geological Survey (USGS; <http://waterdata.usgs.gov/nwis>) has recorded stream flows within the watershed since the early 20th century. Over most of that period, gauged flow of the Muskegon River from USGS, which is the primary input to the estuary, shows an increase of 34% in mean flow, 16% in low flow, and 10% in peak flow since monitoring began in 1935. Wiley et al. (2010) showed an increase in the base flow, storm flow, and median discharge by 15–20% in BAU (end of century with business as usual land management) scenario under the current climate regime.

The MLE was listed as an EPA AOC in 1985 due to historical contamination and habitat degradation. Recent observations in the MLE and previous annual samplings have revealed an annually recurring lake-wide hypoxia (Biddanda, 2012; Biddanda et al., 2018), and the potential for hypolimnetic habitat degradation and increased

eutrophication due to sediment phosphorus release (Steinman et al., 2008; Weinke and Biddanda, 2018).

Preliminary results from observations suggest that hypoxia in the estuary is mitigated by episodic cold-water intrusion from Lake Michigan during coastal upwelling events under certain wind conditions (Biddanda et al., 2018). Water temperature and dissolved oxygen throughout the water column were measured by the Muskegon Lake Observatory (MLO; Fig. 1a) each year from April/May to November/December since 2011 (Biddanda, 2012; Biddanda et al., 2018; [www.gvsu.edu/buoy](http://www.gvsu.edu/buoy)). Physically, MLO observations revealed that the cold-water intrusion strengthens MLE's stratification in the middle of July, August, and early September as shown in Fig. S3a. Ecologically, the dissolved oxygen saturation (Fig. S3b) showed that the hypoxia is mitigated by the cold-water intrusion with increased dissolved oxygen concentrations during the intrusion.


The cold-water intrusion's influence on the ecosystem is more complex than what is revealed from a single station due to the spatial variations in nutrients and seston from the Muskegon River to Lake Michigan. Observations at five fixed stations in MLE and the near shore zone of Lake Michigan from March to October 2003 exhibited explicit spatial variations in nutrient concentrations and seston stoichiometry from the Muskegon River to Lake Michigan (Marko et al., 2013). Much of the terrestrial material from the Muskegon River was intercepted and processed by MLE before flowing into Lake Michigan. Therefore, generally, levels of organic matter (including dissolved organic carbon and particulate organic carbon), total suspended matter, and total phosphorus contained in the fine-grain sediment is consistently lower in Lake Michigan than those observed in MLE (Marko et al., 2013). Stratification combined with the high amount of organic matter in MLE leads to episodic hypoxia in the estuary (Biddanda et al., 2018), while such an event is infrequent in Lake Michigan. As another major spatial difference, because Lake Michigan has phosphorus limitation to phytoplankton growth (Hecky et al., 1993), the mean C:P ratio was much higher in Lake Michigan than MLE due to the lower phosphorus concentrations in Lake Michigan. Thus, water exchange between the Great Lakes and freshwater estuaries is closely associated with nutrients/organic matter redistribution.

In light of the significance of cold-water intrusions to water exchange between the Great Lakes and freshwater estuaries, modeling of cold-water intrusions and its response to different factors is valuable for the understanding of estuarine ecosystems. In this work, we investigate the processes that govern exchange in a freshwater estuary. Through the combination of a long record of observations in the MLE and the development of a hydrodynamic model, we explore the dynamics behind dense water intrusion to freshwater estuaries during coastal upwelling, as well as the intrusion's response to shifts in hydrologic inputs that would result under changing precipitation patterns as a result of climate change. Finally, we test the sensitivity of the dense water intrusion to anthropogenic activities such as alterations to the navigation channel that connects the estuary to the coastal waters.

## 3. Methods

### 3.1. Model description

To understand the physical processes that govern exchange in the MLE, a three-dimensional hydrodynamic model has been developed for MLE, which extends from the mouth of the Muskegon River to the offshore of Lake Michigan (Fig. 1). The model is based on the Semi-implicit Cross-scale Hydroscience Integrated System Model (SCHISM; Zhang et al., 2011, 2015, 2016), an open-source community-supported modeling system based on unstructured grids, derived from the early SELFE model (Zhang and Baptista, 2008). It employs a highly efficient and accurate semi-implicit finite-element/finite-volume method with Eulerian-Lagrangian algorithm to solve the Navier-Stokes equations, and has been widely applied to bays and estuaries around the world



**Fig. 1.** a) Bathymetry of Muskegon Lake Ecosystem (MLE) with observing buoy locations, and (b) coastal Lake Michigan with Semi-implicit Cross-scale Hydroscience Integrated System Model (SCHISM) configuration.

(Zhang et al., 2015; Liu et al., 2017; Chao et al., 2017). The model uses the two-equation closure schemes from the GLS model of Umlauf and Burchard (2003) with the stability function by Kantha and Clayson (1994), and a quadratic bottom drag formulation for bottom stress. The model domain includes Muskegon Lake and the eastern coast of Lake Michigan at the mouth of MLE (Fig. 1b) with horizontal resolution ranging from approximately 2 km in the coastal area in Lake Michigan to as high as 25 m in the navigation channel and MLE, and vertical resolution defined by 20 terrain-following sigma layers. Model bathymetry and coastline topography were provided by NOAA global relief ([ngdc.noaa.gov](http://ngdc.noaa.gov)) and Electronic Navigation Charts ([encdirect.noaa.gov](http://encdirect.noaa.gov)).

Unlike in most coastal ocean systems, tides in the Great Lakes are

weak, and thus the dominant mechanism for transport is the surface wind field, which produces episodic circulation patterns, storm surges, and seiches (Mortimer, 2004). In Lake Michigan, long-term observations found cyclonic circulations in summer and increased cyclonic circulation in winter, important for the transport pathways of nutrients and contaminants on long time scales (Beletsky et al., 1999). In order to account for the complex hydrodynamics in Lake Michigan and the impact on the estuary, the MLE model uses a one-way nesting approach from the NOAA Lake Michigan-Huron Operational Forecast System (LMHOFS; Anderson and Schwab, 2017; <https://tidesandcurrents.noaa.gov/ofs/glofs.html>).

Hourly surface forcing is prescribed from the Great Lakes Coastal

Forecasting System (GLCFS; Schwab and Bedford, 1994), which provides spatially varying wind, air temperature, humidity, long wave and short wave radiation. The GLCFS uses real-time coastal observations from around the Great Lakes to generate a spatially-interpolated meteorological field that is accurate for over-lake conditions. The major hydrologic input to the MLE is from the Muskegon River, which drains the second largest watershed in Michigan and connects at the eastern side of the lake. Hourly river discharge from the USGS Croton Dam station, which is located 52 km upstream from the river mouth, is scaled using an area-ratio method of the total and the gauged-portion of the watershed.

### 3.2. Numerical experiments design

A control case simulation was carried out for the period January 1 – December 31, 2016 using lateral and surface boundary conditions as described above, and initialized with a constant water temperature at 2 °C. This period covers the formation of thermal stratification in the early summer, several upwelling-induced cold-water intrusion events during the stratified period, and thermal mixing and overturning in the fall. Modeled thermal structure and currents were compared to observations within the MLE at CHAN, WEST, DEEP and EAST sites (Fig. 1a) and along the shore of Lake Michigan at NOAA station 45161 (Fig. 1b). Observations at the MLE sites were sampled every hour for water quality parameters and temperature; at NOAA station 45161, temperature and currents are sampled every 20 min. In addition, a series of idealized experiments were carried out to examine the response of cold-water intrusion to different factors (Table 1). The idealized experiments were focused on a period of significant cold-water intrusion that occurred in September. In each case, the model was restarted from August 28 using initial conditions from the control case simulation.

To examine the response of cold-water intrusion to shifts in hydrologic inputs under changing precipitation patterns (Wang et al., 2017), in experiments R1, R2, R3 and R4, the variations in river discharge were simulated by multiplying the flow from August 28 to December 31 by 0.5, 2, 3 and 4 times with averaged flow rates during the intrusion of 20, 80, 120 and 160 m<sup>3</sup>/s, respectively. Fig. S2 shows the historical discharge observations from 1996 to 2016, demonstrating the inter-annual variability of hydrologic input in summer when the water in Lake Michigan and the MLE is stratified. While climate projections suggest intensification and shift of extreme precipitation events over the next century (Donat et al., 2016; Wang et al., 2017), the cases R3 and R4 are used to examine the change of cold-water intrusion under climate projection including intensification and shift of extreme

precipitation events.

Anthropogenic activities may affect the coastal ocean, estuaries and lakes with many stressors including urbanization and runoff, agriculture, microbial processing of organic matter and nitrification/denitrification (Hoffman et al., 2014), as well as shoreline modifications such as shoreline engineering, dune destruction, channel dredging and jetty construction. Even though the navigation channel is unlikely to change in MLE, the influence of navigational channel characteristics on hydrodynamic and ecological processes is relevant to extending this work to other similar systems. In addition, previous studies have suggested that the potential for Lake Michigan waters to flow into MLE is limited due to the narrow connection in the channel (Marko et al., 2013), and thus our work aims to verify those conclusions. In experiments N1, N2 and N3, the impact of the navigation channel was evaluated by broadening its width by 2, 3, and 4 times, respectively.

Finally, the strength of upwelling and consequent extent of cold-water intrusion depends on the wind stress and nearshore bathymetry. To investigate the intrusion's response to upwelling strength under prevailing alongshore winds with different magnitudes (Table 1), experiments W1–4 are undertaken. In each case, prevailing alongshore upwelling-favorable winds are applied for the period August 28–30 at 3 m/s, 5 m/s, 7 m/s and 15 m/s, respectively, followed by no wind for 10 days.

## 4. Results and discussion

### 4.1. Model assessment

Before analyzing model results from the idealized experiments, the control case simulation was first validated by comparing with available observations to establish confidence in the model. To verify the modeled stratification and currents in Lake Michigan, comparisons were made with measured temperature and velocities at NOAA station 45161 (Fig. 2a and b). Currents are observed at NOAA 45161 from 2.5 m to 20.5 m depth with a bin of 1 m. We compared the daily-averaged, model currents with the observations (Fig. 2a) at selected depths throughout the water column.

According to Fig. 2a, the modeled currents are consistent with observations in amplitude and change of direction with root mean square errors (RMSEs) of 5.30, 4.44, 3.77, 3.03 and 2.53 cm/s for u-component velocity at 2.5, 7.5, 12.5, 17.5 and 20.5 m below the surface, and 10.5, 9.39, 7.09, 4.40 and 3.12 cm/s for v-component (Table 2). When upwelling-favorable winds occur during the beginning of August and September, the model captures the strong surface currents moving southeastward. The currents decrease with depth and reverse at a depth. For example, the current reverses at the depth of 12.5 m at the beginning of August. To further evaluate model performance and the hydrodynamics in the nearshore, additional statistical measures are included to be consistent with previous studies of Lake Michigan (Eqs. (1)–(4); Schwab, 1983; Beletsky and Schwab, 2001; Anderson and Schwab, 2011, Table 2).

$$\sigma_o = \frac{\sum [(u_o - \bar{u}_o)^2 + (v_o - \bar{v}_o)^2]}{\sum (u_o^2 + v_o^2)} \quad (1)$$

$$\sigma_c = \frac{\sum [(u_c - \bar{u}_c)^2 + (v_c - \bar{v}_c)^2]}{\sum (u_c^2 + v_c^2)} \quad (2)$$

$$\varepsilon = \frac{\sum [(u_o - \bar{u}_o - u_c + \bar{u}_c)^2 + (v_o - \bar{v}_o - v_c + \bar{v}_c)^2]}{\sum [(u_o - \bar{u}_o)^2 + (v_o - \bar{v}_o)^2]} \quad (3)$$

$$v_o, v_c = \left( \frac{1}{n} \sum_{i=1}^n |v_o - v_c|^2 \right)^{1/2} \quad F_n = \frac{v_o, v_c}{v_o, 0} \quad (4)$$


where ( $u_o$ ,  $v_o$ ) represent observed current, and ( $u_c$ ,  $v_c$ ) represent modeled (computed) current.

**Table 1**  
Process-oriented experiments for the study of intrusion length.

Run	Width of Navigation Channel	River Discharge	Wind Speed
Ctrl case	Real	Real	Real
R1	Real	Half	Real
R2	Real	2 times	Real
R3	Real	3 times	Real
R4	Real	4 times	Real
N1	2 times	Real	Real
N2	3 times	Real	Real
W1	Real	Real	Alongshore winds; 3 m/s <sup>a</sup>
W2	Real	Real	Alongshore winds; 5 m/s
W3	Real	Real	Alongshore winds; 7 m/s
W4	Real	Real	Alongshore winds; 15 m/s

<sup>a</sup> The alongshore winds in W1, W2, W3 and W4 are upwelling favorable, i.e., northwesterly, for the first three days, after three days, there is no wind applied.





**Fig. 2.** Comparisons of currents and temperature between model and observations including a) currents at station NOAA 45161 in Lake Michigan at 2.5, 7.5, 12.5, 17.5 and 20.5 m below surface, surface and bottom temperature b) at NOAA 45161, c) at station CHAN inside the navigation channel, and d) at station DEEP. The arrow direction in a) represents the current direction with an arrow pointing northward for a northward current.

**Table 2**  
Comparisons of modeled and observed currents at NOAA 45161.

Depth (m)	RMSE_u (cm/s)	RMSE_v (cm/s)	$\sigma_o$	$\sigma_c$	$\varepsilon$	$F_u$	$F_v$
2.5	5.30	10.50	0.94	0.97	0.49	0.69	0.70
7.5	4.44	9.39	0.96	1.00	0.51	0.70	0.75
12.5	3.77	7.09	1.00	0.97	0.50	0.70	0.73
17.5	3.03	4.40	0.94	0.95	0.49	0.68	0.68
20.5	2.53	3.12	0.84	0.97	0.49	0.66	0.74

$\sigma_o$  ( $\sigma_c$ ) is the ratio of energy in the time-variable component of the currents to the total current energy for observed (modeled) currents. With values from 0.84 to 1.00 at NOAA 45161 (Table 2), it describes the nearshore of Lake Michigan as wind-dominated with a higher ratio of energy in the time-variable component of the currents to the total current energy.

The differences between observed and modeled currents for the time-variable flow are measured by  $\varepsilon$ . A value of 0.0 indicates a perfect match between the observed and modeled currents, and a value of 1.0 indicates an error in the modeled current as large as the observed current. According to previous studies (Schwab, 1983; Schwab and Bennett, 1987; Anderson and Schwab, 2011), a value 0.9 is typical for the time-variable component in the Great Lakes, as the observed currents can contain substantial energy beyond the model resolution. Therefore, prediction with  $\varepsilon \leq 0.9$  is considered successful.


$F_n$ , a normalized Fourier norm, describes the relative percentage of variance in the observed currents that is unexplained by the modeled currents. If  $F_n = 0$ , the model has perfect prediction; If  $0 < F_n < 1$ , the model prediction is better than no prediction at all. For a reference, Schwab (1983) got  $0.79 < F_n < 1.01$  for a barotropic simulation of Lake Michigan currents on a 5-km grid. Beletsky and Schwab (2001) calculated  $0.5 < F_n < 0.9$  for the Lake Michigan coastal current prediction in winter also on a 5-km grid.

At NOAA 45161, with  $\varepsilon$  around 0.5 throughout the water column, and both  $F_u$  ( $F_n$  for the u-component velocity) and  $F_v$  ( $F_n$  for the v-component velocity) around 0.7, the model is found to predict

nearshore currents in Lake Michigan on par with previous studies.

Fig. 2b shows the comparison of surface and bottom temperature at NOAA 45161 during May 2016 to November 2016 with RMSEs of 1.4 °C and 3.1 °C respectively. The surface temperature captures the seasonal trend shown in the observations, and the major events, such as the episodic, abrupt decrease in temperature around June 6, July 2, August 3, and September 2. The model slightly overestimates the bottom temperature, which may arise from a diffuse resolution of the thermocline from the LMHOFs model or from the lack of groundwater flux in the model, which is more obvious in the comparisons of bottom temperature inside MLE.

With respect to conditions inside the MLE, Fig. 2c shows the comparisons of surface and bottom temperature at the station CHAN with RMSEs of 0.6 °C and 2.8 °C, respectively. In particular, the bottom temperature in the channel reveals the model's ability to capture the intrusion events driven by upwelling in Lake Michigan. However, further inside the MLE, it is a more complicated situation. The model's surface temperature closely follows the observations at the DEEP station, with RMSE of 0.7 °C, however, the bottom temperature is highly overestimated (Fig. 2d) with a RMSE of 5.8 °C (similar to other stations in MLE; Fig. S4 in the Supplementary Material). A possible explanation for the discrepancy in bottom water temperature within the MLE could be the inclusion of groundwater inflow to the system. In a similar nearby system, Gull Lake, Safaie et al. (2017) showed groundwater input to be a critical feature to capturing the thermal stratification. Thus, in a simulation of the control case with groundwater inflow included, based on rough approximations of flux and temperature, the model reveals a better resolution of bottom water temperatures within the MLE (Figs. S7–S9 in the Supplementary Material), suggesting that this might be a reasonable culprit and worthy of further investigation. However, in the absence of accurate groundwater information, and since the difference in model results with and without groundwater included are small or negligible in terms of the model's ability to resolve water exchange and the cold-water intrusions driven by upwelling in Lake Michigan, the following model simulations are shown for the control case without groundwater. Further discussion can be found in



**Fig. 3.** a) Daily averaged surface temperature from August 28, 2016 to September 14, 2016 during an upwelling event from the LMHOFS model; b) Winds in MLE in September 2016 with red arrow represents the data used by GLCFS and blue arrow represents the observations at the GVSU buoy site (<http://www.gvsu.edu/wri/buoy/>); c) observed near surface and bottom temperature in September 2016 at a station inside the navigation channel.

section 4.7 and in the Supplementary Material.

#### 4.2. Cold-water intrusion


Along the eastern boundary of Lake Michigan, southward along-shore winds induce Ekman transport (Ekman, 1905) of surface water away from the shore. These displaced surface waters are replaced by deeper Lake Michigan waters, resulting in coastal upwelling often characterized by cold surface waters at the coast (Plattner et al., 2006). Surface temperature from the LMHOFS model in 2016 (Fig. 3a) reveals periodic occurrence of upwelling along the eastern shore of Lake Michigan with surface temperature decreasing to below 18 °C on September 2 and 4, when the lake is stratified, and upwelling-favorable winds develop (Fig. 3b). Again, on September 11, northerly winds occur, though with a much shorter duration than the event beginning on September 1 (Fig. 3b). Correspondingly, there is a weaker upwelling with surface temperature around 20.5 °C (Fig. 3a). Historical observations inside the navigation channel between Lake Michigan and MLE

show colder bottom temperatures during these upwellings events (Fig. 3c).

Taking the large upwelling in September as a representative event to investigate the cold-water intrusion in MLE and its effects to the hydrodynamics of MLE, we plotted the daily averaged bottom temperature in MLE from August 31, 2016 to September 10, 2016 (Fig. 4a–f). Model simulations of temperature and along-channel velocity at CHAN (Fig. 4 g–i) demonstrate that the episodic occurrence of bottom cold-water in Figs. 2 and 3 is due to the cold-water intrusion from Lake Michigan to Muskegon Lake during upwelling-favorable winds (Fig. 3b).

On August 28 and 29, before the upwelling, the water column at the NOAA station 45161 was highly stratified with surface temperature higher than 20 °C (Fig. 4g). The subsurface water temperature fluctuates with a period of ~17 (about 17.2) h, which is caused by the oscillations of rotation-modified internal (Poincaré) wave (Schwab, 1977; Beletsky et al., 1997). Poincaré waves are a travelling disturbance of the wind-induced upwelling/downwelling perturbations of





**Fig. 4.** SCHISM model results for the cold-water intrusion event at the beginning of September 2016. a-f) Daily-averaged bottom-temperature during the cold water intrusion event; Hovmöller diagrams for f) temperature at NOAA 45161, g) temperature at the station CHAN in the navigation channel and h) velocity along the navigation channel for the station CHAN in the channel. The positive velocities represent currents moving into MLE, and negative ones represent currents out of MLE.

the thermocline on a whole-basin scale (Schwab, 1977; Mortimer, 1984). They are characterized by clockwise phase progression with a period slightly less than the local inertial period,  $2\pi/f$  ( $f$  is Coriolis parameter), 17.5 h for Central Michigan. Spectral analysis of water elevation reveals the oscillation also occurs in the navigation channel and inside MLE, although its influences to the temperature profile in the channel are negligible (Fig. 4h) - the navigation channel is well mixed

with temperature higher than 20 °C through the water column. Before the upwelling, along-channel velocity inside the navigation channel is characterized by a high frequency oscillation with a period of 3.25 h (or 3.4, 3.6 h for different time), which is corresponding to the third longitudinal seiche mode in Lake Michigan (Mortimer, 2004). Spectral analysis of our model results shows the oscillation occur in both Lake Michigan and MLE, but relatively stronger in MLE. However, the seiche

is not strong enough to affect the reversals in water exchange within the channel. Currents in the channel move from MLE to Lake Michigan through the water column with a speed about 10 cm/s (Fig. 4i).

From September 1 to September 7, the Ekman transport moves surface water away from the eastern shore of Lake Michigan, and replaces the surface water with bottom offshore water, decreasing temperature through the water column near shore, for example, at NOAA 45161 from 25 °C to 5 °C (Fig. 4g). During upwelling, the Poincaré wave still dominates the thermocline oscillations in Lake Michigan but with lower temperature. The replaced water moving further into the MLE through the connection channel, stratifies the channel's water column with a cold subsurface layer from Lake Michigan waters, and induces a bi-directional flow in the channel, where cold-waters intrude along the bottom half of the channel. As a result of the cold-water intrusion, the surface current strengthens from MLE to Lake Michigan, although the change in net flow was negligible.

The cold-water intrusion first moves northeastward to the northern deep trough toward the mouth of Bear Creek, and then moves gradually southeastward along the southern trough reaching ~80% of MLE (Fig. 4). Around September 6, when downwelling-favorable winds predominate over upwelling-favorable winds, the upwelling along Lake Michigan's coast retreats, leaving the cold-water in MLE and gradually dissipating by mixing. By calculation, the total volume of the cold-water intruded to MLE during the event from September 1 to September 6 is  $1.57 \times 10^7 \text{ m}^3$ , which is 13% of the Muskegon Lake volume ( $1.25 \times 10^8 \text{ m}^3$  from the SCHSIM model configuration). The net exchange was around  $2.2 \times 10^7 \text{ m}^3$ , close to the total hydrologic input to MLE during this event at an average flow rate of 40 m<sup>3</sup>/s.

Although intrusion doesn't significantly alter the net outflow, it initiates the water exchange between Lake Michigan and MLE by bringing oligotrophic water into MLE from the bottom and facilitating stronger mesotrophic outflow to Lake Michigan from the surface. With the occurrence of cold-water intrusion, fewer nutrients and seston are retained in MLE. According to available studies over MLE (Marko et al., 2013; Weinke et al., 2014), the spatial nutrients/seston distribution from the Muskegon River to Lake Michigan plays a significant role in the system's ecological processes. Therefore, the cold-water intrusion is important for the ecological processes.

#### 4.3. Response to hydrologic input shift

As climate projections reveal intensification of extreme precipitation events over the next century (Donat et al., 2016; Wang et al., 2017), experiments R1 to R4 are used to examine how hydrologic inputs affect cold-water intrusions into the estuary. During the upwelling intrusion at the beginning of September, the averaged flow rates from the Muskegon River for the experiments R1, Ctrl, R2, R3 and R4 are 20, 40, 80, 120, and 160 m<sup>3</sup>/s, respectively. For all these experiments, we released a conservative, passive dye at the bottom of the water column inside the jetties on September 1 to track the cold-water intrusion into the estuary. An instantaneous snapshot of the horizontal distribution of the dye tracer on September 5 shows that with an increase in river discharge, the intrusion length decreases (Fig. 5). When the average river discharge becomes 160 m<sup>3</sup>/s, the cold-water intrusion is completely suppressed.

We quantified the intrusion difference by the change in water volume into MLE across the navigation channel (Fig. 5f). Halving the river discharge increases the transport of bottom water from Lake Michigan to MLE from ~13% of MLE's total volume to ~18%, while increases in river discharge suppress the volume of Lake Michigan waters entering the estuary. For example, doubling the discharge decreases the intrusion volume to ~8%. When the river discharge increases to 4 times of the original value, the intrusion was suppressed with delivery close to 0. The differences between the river input to MLE and the water leaving MLE through the navigation channel was close to the intrusion volume, with a small net balance keeping the total volume in MLE changes

slightly.

#### 4.4. Response to change in navigation channel width

Experiments N1 and N2 are used to demonstrate the influence of connecting channel characteristics on exchange between drowned river-mouth estuaries and offshore waters. Fig. 6 shows the bottom temperature distribution for this set of experiments. With the increase of navigation channel width, the intrusion length increases. For the case Ctrl (Fig. 6a), waters colder than 15 °C stays in the northern trough only, while in the case N2, they reach 1/3 of the total MLE's length.

The intrusion difference was also quantified based on the change in volume flux from Lake Michigan to MLE as shown in Fig. 6d. Compared with the base experiment, doubling the navigation channel width increases the intrusion flux from 13% of Muskegon Lake's volume to 20%, and to 22% if tripling the width. The change of outflow from MLE is proportional to the change in cold-water intrusion, resulting in a small change in the net balance. Despite that, broadening the channel significantly increases the water exchange between Lake Michigan and MLE. This study will provide a good reference for the future studies of other similar brackish or freshwater estuaries that are connected to larger open water bodies with a channel.

#### 4.5. Response to wind conditions


Wind experiments W1 to W3 have wind speeds that are common in Lake Michigan, while W4 has an extreme wind speed as shown in Fig. S5. The Hovmöller diagrams in Fig. 7a–h show the velocity and temperature in the navigation channel. We can see the increase of wind amplitude from 3 m/s in W1 to 7 m/s in W3 strengthens the intrusion with stronger vertical shear in currents and larger temperature stratification. The intrusion flux (Fig. 7i) increased from about 10 m<sup>3</sup>/s in W1 to 20 m<sup>3</sup>/s in W3, while the net flux to MLE keeps nearly constant. Meanwhile, the onset time of intrusion becomes earlier from W1 to W3 with the increase of wind speed, which resulted from the quick build-up of pressure gradient between offshore and onshore regions. Fig. 7j shows the pressure gradient between two stations on the near shore region of Lake Michigan (ends of the magenta line in Fig. 1b). For W1–3, it increases gradually during the first three days with winds, and then decreases smoothly after winds' disappearance. The intrusion can continue for another several days, dependent on the time needed for pressure restoration. Stronger upwelling generated a longer tail for the intrusion flux; meanwhile the cold-water moved into MLE with a larger intrusion length, i.e., the length that cold-water propagates in MLE (Fig. 8). In W1, the cold-water intrusion was weak and difficult to identify, though in W2 and W3, the intrusion length increased in response to the increase in wind forcing.

Extreme winds can occur along the eastern coast of Lake Michigan, affecting its estuaries' hydrodynamics and ecosystem. For example, when Hurricane Sandy's remnants reached Lake Michigan in October 2012, wind gusts were as strong as 17 m/s. Using an experiment with an extreme wind events for three days, we investigated how the potential extratropical storms affect MLE's cold-water intrusion.

In W4, where the applied wind is akin to a northwesterly, along-shore extratropical storm, upwelling along the eastern shore of Lake Michigan is strong. However, due to the strong turbulent mixing in the navigation channel and MLE, during the applied wind storm, the stratification and vertical shear in the channel (Fig. 7g–h) and MLE (Fig. 8) were suppressed. Therefore, the intrusion flux through the channel was weak, even though the pressure gradient perpendicular to the shore is strong. After the removal of winds, the forcing that balances the strong pressure gradient disappear, which drives a strong transient current toward MLE. With the rapid decrease of pressure gradient and the weakening of turbulent mixing, the cold-water intrusion occurred after the removal of winds during the restoration of pressure gradient.

Another form of upwelling-favorable wind, offshore wind blowing






**Fig. 5.** Model results for the experiments with different hydrologic input during the major upwelling on September 05, 2016 12:00:00, including passive tracer distributions for the experiments a) R1, b) Ctrl, c) R2, d) R3 and e) R4 with normalized river discharges of 0.5, 1, 2, 3 and 4, respectively, and f) the corresponding changes in water exchange.

from the MLE to Lake Michigan, also causes upwelling along the eastern shore of Lake Michigan and cold-water intrusion into the MLE. Because the change of wind speed in offshore wind causes the similar response in cold-water intrusion by the change in alongshore wind, we put the related model results (Fig. S6) and discussion in the Supplementary Material.

#### 4.6. Frequency of cold-water intrusions

Continuous hydrodynamic monitoring in the navigation channel

between MLE and Lake Michigan was not established until 2016. Therefore, direct measurements of cold-water intrusion were not available before 2016. However, as a direct consequence of upwelling along the eastern shore of Lake Michigan, the cold-water intrusion can be inferred by the observation of upwelling at coastal locations. When stratification is established in Lake Michigan, upwelling decreases temperature through the water column; in contrast, downwelling piles up warm surface waters toward the coast, increasing deep water temperature. Buoys deployed in the nearshore Lake Michigan have been recording the near-surface and near-bottom temperature since 2009



**Fig. 6.** Bottom temperature on September 05, 2016 12:00:00 for the experiments a) Ctrl, b) N1 and c) N2 with normalized navigation channel widths of 1, 2 and 3, respectively, and d) their corresponding changes in water exchange. In N1 and N2, the navigation channel in Ctrl experiment is broadened by 2 and 3 times, respectively.

(NOAA's National Data Buoy Center Station 45161; <http://www.ndbc.noaa.gov/>). Therefore, we could analyze the temperature anomalies to identify upwelling along the eastern shore of Lake Michigan and, hence, the cold-water intrusion to MLE.

Fig. 9 shows the near-surface (dashed lines) and near-bottom (solid lines) temperature anomalies, the temperature variation by removing seasonal trend, at station 45161 from 2015 to 2017. An upwelling event is established accompanied by the decrease of both near-bottom and near-surface temperature. When the upwelling doesn't have enough duration or strength, the near-surface temperature anomaly is small. We marked the potential upwelling-caused water anomalies as upward arrows in Fig. 9. The intrusions were likely to occur at least once from June to November in 2015 and 2016, and likely more prevalent in 2017. The temperature anomalies in July 2017 show 3 upwelling events spanning the whole month. The duration of cold-water intrusion is mostly determined by the upwelling duration according to the previous analysis. Fig. 9 shows that the cold-water intrusion can last as long as weeks. For example, the intrusion in August 2016 lasted about 3 weeks, and the intrusion at the beginning of September 2016 lasted for a week. However, there is risk to counting the intrusion duration solely by the upwelling duration, because the intrusion is susceptible to the river flow suppression and strong wind mixing in the channel.

#### 4.7. Model limitations

Groundwater is another important natural input to the Great Lakes, accounting for 1–12% of total water inflow to Lake Michigan (Robinson, 2015), as well as to the freshwater estuary of MLE along Lake Michigan. However, due to the statistical variability and methodological inaccuracies associated with the monitoring efforts, groundwater data is scarce and imprecise (Mitchell et al., 1988). To examine the impact of groundwater flux to the hydrodynamics of MLE, we arbitrarily added 20 groundwater sources along the deepest locations in the eastern reach of the MLE (marked in Fig. S7) with a total constant flux of  $20 \text{ m}^3/\text{s}$  and temperature of  $8^\circ\text{C}$ . According to Collins (1925), the approximate average temperature of shallow groundwater obtained from shallow wells in western Michigan ranges from  $8^\circ\text{C}$  to  $11^\circ\text{C}$ . In the idealized experiment, we used even lower temperature for the highly sparse groundwater sources. The modeled temperature (Fig. S8) shows significant improvement in bottom temperature and stratification. The idealized experiment demonstrates the groundwater flux could significantly increase MLE's stratification. Although the groundwater input may play an important role in the MLE's ecosystem, as a potential source of nutrients that stimulates algal blooms (Kilroy and Coxon, 2005; Lewandowski et al., 2015; Knights et al., 2017), its effect on timing and duration of cold-water intrusion is small (Fig. S9). However, it can slightly decrease the volume of the intrusion (around 10%) according to the time series of cold-water intrusion flux (Fig. S10).





Fig. 7. Water exchange between Lake Michigan and MLE from the experiments with prevailing upwelling-favorable winds in different magnitudes of 3, 5, 7 and 15 m/s in W1 to W4. a-h) Hovmöller diagrams for along-channel currents and temperature at the CHAN station for experiments W1 to W4; i) Intrusion flux into MLE (solid lines) and net flux into MLE (dashed lines) for the experiments W1 to W4; j) Pressure gradients between the onshore and offshore points in Lake Michigan close to MLE. The onshore and offshore points in Lake Michigan are the ends of the magenta line shown in Fig. 1b.





**Fig. 8.** Bottom temperature snapshots on August 30, 2016 12:00:00 for experiments with prevailing upwelling-favorable winds from August 28 to August 30 in a) 3 m/s (W1), b) 5 m/s (W2), c) 7 m/s (W3) and d) 15 m/s (W4).

Further improvement of the SCHISM model with realistic groundwater sources will be conducted in the future.

## 5. Summary and conclusions

We used a 3-D hydrodynamic model (SCHISM) to study the water exchange between Lake Michigan and a freshwater estuary. The model successfully captures the episodic cold-water intrusions driving by upwelling events in Lake Michigan. As a result of Ekman transport, the upwellings deliver a large volume of cold-water to MLE, resulting in a significant impact on the hydrodynamics of the freshwater estuary. For example, the upwelling at the beginning of September 2016 induced a cold-water intrusion with a volume near 13% of the entire MLE. For the first time, near-surface and near-bottom temperature anomalies at NDBC 45161, a NOAA buoy along Lake Michigan's eastern shore, are analyzed to indirectly identify the occurrence of cold-water intrusion from Lake Michigan to MLE. The results reveal that the cold-water intrusion to MLE is common, which appears a least once per month with duration from days to weeks, determined by the duration and strength of the upwelling-favorable winds.

In addition to hydrodynamic implications, these intrusions exchange biologically distinct waters between Lake Michigan and MLE. Compared with MLE, waters in Lake Michigan contain less organic matter, total suspended matter, and total phosphorus of fine sediment, but more dissolved oxygen. To conserve the volume in MLE, the cold-water intrusion from Lake Michigan to MLE along the lake bottom is accompanied by a stronger surface outflow. Therefore, the intrusion not

only dilutes the related matter and nutrients in MLE but also accelerates their transport out of MLE. Furthermore, [Biddanda et al., \(2018\)](#) suggested that the episodic cold-water intrusion affects the MLE's ecosystem by bringing water with more dissolved oxygen to the estuary bottom, mitigating summer hypoxia for a short period of time, in contrast to the commonly-observed intrusion of hypoxic waters from coastal oceans into brackish estuaries ([Deppe et al., 2017](#); [Rabalais et al., 2002](#); [Breitburg, 1992, 2002](#); [Sanford et al., 1990](#); [Levin et al., 2009](#)).

Through process-oriented experiments, we examined the cold-water intrusion's responses to hydrological shift, variation in navigation channel width, and wind conditions. It is found that the increase in riverine input during the upwelling weakens the intrusion. [Marko et al. \(2013\)](#) suggested the increase in river discharge brings more organic matter, total suspended matter, total phosphorus and sediment into MLE, while our study demonstrates the suppression of the intrusion by the increase in river discharge. The suppression may deteriorate the water quality by retaining more organic matter and nutrients in MLE – including summer time surface blooms of harmful cyanobacteria. This would suggest that a hydrologic shift under a changing climate would alter the ecosystem through affecting the water exchange between Lake Michigan and MLE. The conclusion may provide a good reference to the study of estuaries whose physical and ecological processes are susceptible to upwelling-downwelling wind cycle and changing hydrological conditions, for example, San Francisco Bay ([Liu et al., 2018](#)), Chesapeake Bay ([Breitburg, 1992, 2002](#)), the Columbia River Estuary ([Roegner et al., 2011](#)), Willapa Bay ([Roegner et al., 2002](#)), etc.




Fig. 9. Anomalies of surface temperature and bottom temperature measured at the NOAA 45161 station in a) 2015, b) 2016 and c) 2017. A high-pass filter with a cut-off period of 1 month was used to remove the seasonal trend. Arrows indicate periods of potential upwelling.

In addition, the present work shows that the extent and strength of intrusion depends on the structure of the connecting pathways between the estuary and the offshore waters. Results show that the increase of navigation channel width strengthens the cold-water intrusion, which is not only important for extrapolation of these results to other freshwater estuaries, but may also be important for management and coastal engineering projects in similar systems with narrow channels (for example, Tampa Bay, [Zhu et al., 2014](#)).

Through experiments with different wind speeds in the along-shelf direction, we further investigated the dynamics behind the intrusion. Stronger upwelling causes a stronger and longer duration intrusion with larger intrusion length into MLE. Driven by the pressure gradient between onshore and offshore, the cold-water intrusion doesn't disappear right after the absence of upwelling-favorable winds, but instead dissipates gradually with the relaxation of the pressure gradient. When extreme upwelling-favorable winds occur (for example, during extratropical storms, winds can reach 15 m/s), the intrusion during the winds is weak. However, the strong upwelling caused by the extreme winds continues to work after the winds absence with an even stronger intrusion to MLE.

Overall, cold-water intrusions may be critical for optimal water exchange and ecosystem function of temperate freshwater estuaries. Such hydrodynamic connectivity may be especially relevant in systems such as the Great Lakes or in the world's coastal estuaries where tidal forcing is weak, including White Lake, Pentwater Lake and Lake Macatawa along the eastern shore of Lake Michigan, Hamilton Harbour ([Lawrence et al., 2004](#)), Frechman's Bay, and White Harbour along Lake Ontario, etc. The frequency and extent of these intrusions depends on a

number of factors. However, the present work suggests that under shifting conditions, such as winds and precipitation events or intensity, the critical delivery of nutrients and other water quality components can be altered dramatically, potentially eliminating the replenishment of high dissolved-oxygen and low nutrient intrusions and exacerbating both hypoxic and eutrophic conditions in estuaries.

#### Acknowledgements:

We thank the NOAA-Great Lakes Environmental Research Laboratory (GLERL)'s Lake Michigan Field Station for providing logistical ship-support with yearly Muskegon Lake Observatory buoy (MLO) deployment and recovery operations. The MLO infrastructure and open access data that were utilized in this study was funded by EPA Great Lakes Restoration Initiative (R5-GL2010-1), NASA Michigan Space Grants Consortium (NNX15AJ2018) and University of Michigan-Cooperative Institute for Great Lakes Research (CIGLR) grants (NA12OAR4320071) to BB. We are grateful to Steve Ruberg, NOAA-GLERL for open access data from NOAA's nearshore Lake Michigan buoy (NOAA 45161). A Community Foundation for Muskegon County grant to the Annis Water Resources Institute supported the long-term seasonal ship-board monitoring of Muskegon Lake. We thank Kurt Thompson and Alan Steinman, Annis Water Resources Institute, Grand Valley State University for providing regional groundwater data for analysis. QL acknowledges the support from a CIGLR postdoctoral fellowship award (NA12OAR4320071) to BB and EA and the CIGLR grant through the NOAA cooperative agreement NA17OAR4320152. This is CIGLR publication #1128 and GLERL publication #1886.

## Appendix A. Supplementary data

Supplementary data related to this article can be found at <http://dx.doi.org/10.1016/j.ecss.2018.05.014>.

## References

- Anderson, E.J., Schwab, D.J., 2011. Relationships between wind-driven and hydraulic flow in Lake St. Clair and the St. Clair river delta. *J. Great Lake Res.* 37, 147–158.
- Anderson, E.J., Schwab, D.J., 2017. Meteorological influence on summertime baroclinic exchange in the Straits of Mackinac. *J. Geophys. Res. Oceans* 122 (3), 2171–2182.
- Allan, J.D., McIntyre, P.B., Smith, S.D.P., Halpern, B.S., Boyer, G.L., Buchsbaum, A., Burton Jr., G.A., Campbell, L.M., Chadderton, W.L., Ciborowski, J.J.H., Doran, P.J., Eder, T., Infante, D.M., Johnson, L.B., Joseph, C.A., Marino, A.L., Prusevich, A., Read, J.G., Rose, J.B., Rutherford, E.S., Sowa, S.P., Steinman, A.D., 2012. Joint analysis of stressors and ecosystem services to enhance restoration effectiveness. *Proc. Natl. Acad. Sci. U.S.A.* 110, 372–377.
- Borja, Á., Dauer, D.M., Elliott, Simenstad, C.A., 2010. Medium- and long-term recovery of estuarine and coastal ecosystems: patterns, rates and restoration effectiveness. *Estuar. Coast* 33, 1249–1260.
- Biddanda, B.A., 2012. Lake Sentinel: Observatory for Ecosystem Changes in Muskegon Lake. InterChange, Newsletter of Regional Math and Science Center. Grand Valley State University. <http://www.gvsu.edu/rmsc/interchange/2012-may-connections-awri-651.htm>.
- Biddanda, B.A., Weinke, A.D., Kendall, S.T., Gereaux, L.C., Holcomb, T.M., Snider, M.J., Dila, D.K., Long, S.A., VandenBerg, C., Knapp, K., Koopmans, D.J., Thompson, K., Vail, J.H., Ogdahl, M.E., Liu, Q., Johengen, T.J., Anderson, E.J., Ruberg, S.A., 2018. Chronicles of Hypoxia: time-series buoy observations reveal annually recurring seasonal basin-wide hypoxia in Muskegon Lake – a Great Lakes estuary. *J. Great Lake Res.* 44 (2), 219–229. <https://doi.org/10.1016/j.jglr.2017.12.008>.
- Biddanda, B.A., 2017. Global significance of the changing freshwater carbon cycle. *Eos. American Geophysical Union* 98 (6), 15–17. <https://doi.org/10.1029/2017EO069751>.
- Beletsky, D., O'Connor, W.P., Schwab, D.J., Dietrich, D.E., 1997. Numerical simulation of internal kelvin waves and coastal upwelling fronts. *J. Phys. Oceanogr.* 27, 1197–1215.
- Beletsky, D., Saylor, J.H., Schwab, D.J., 1999. Mean circulation in the great lakes. *J. Great Lake Res.* 25 (1), 78–93.
- Beletsky, D., Schwab, D.J., 2001. Modeling circulation and thermal structure in Lake Michigan: annual cycle and interannual variability. *J. Geophys. Res.* 106, 19745–19771.
- Breitbart, D.L., 1992. Episodic hypoxia in the Chesapeake Bay: interacting effects of recruitment, behavior and a physical disturbance. *Ecol. Monogr.* 62, 525–546.
- Breitbart, D.L., 2002. Effects of hypoxia, and the balance between hypoxia and enrichment, on coastal fishes and fisheries. *Estuaries* 25 (767). <https://doi.org/10.1007/BF02804904>.
- Chao, Y., Farrara, J.D., Zhang, H., Zhang, Y.J., Ateljevich, E., Chai, F., Davis, C.O., Dugdale, R., Wilkerson, F., 2017. Development, implementation, and validation of a modeling system for the San Francisco Bay and estuary. *Estuar. Coast Shelf Sci.* <https://doi.org/10.1016/j.ecss.2017.06.005>.
- Chen, S.-N., Geyer, W.R., Ralston, D.K., Lerczak, J.A., 2012. Estuarine exchange flow quantified with isohaline coordinates: contrasting long and short estuaries. *J. Phys. Oceanogr.* 42, 748–763. <http://dx.doi.org/10.1175/JPO-D-11-086.1>.
- Cole, J.J., Prairie, Y.T., Caraco, N.F., McDowell, W.H., Tranvik, L.J., Striegl, R.G., Duarte, C.M., Kortelainen, P., Downing, J.A., Middelburg, J.J., Melack, J., 2007. Ecosystems 10 (1), 172–185. <https://doi.org/10.1007/s10021-006-9013-8>.
- Collins, W.D., 1925. Temperature of water available for industrial use in the United States 1923–1924. In: *Contributions to the Hydrology of the United States, Water Supply Paper 520, Chapter F. United States Geological Survey, Washington, DC*, pp. 97–104.
- Cotner, J.B., Weinke, A.D., Biddanda, B.A., 2017. Great Lakes: science can keep them great. *J. Great Lake Res.* 43, 916–919. <https://doi.org/10.1016/j.jglr.2017.07.002>.
- Deppe, R.W., Thomson, J., Polagye, B., et al., 2017. Predicting deep water intrusions to Puget sound, wa (USA), and the seasonal modulation of dissolved oxygen. *Estuar. Coast* 1–14. <https://doi.org/10.1007/s12237-017-0274-6>.
- Donat, M.G., Lowry, A.L., Alexander, L.V., O'Gorman, P.A., Maher, N., 2016. More extreme precipitation in the world's dry and wet regions. *Nat. Clim. Change* 6, 508–513. <http://dx.doi.org/10.1038/nclimate2941>.
- Ekman, V.W., 1905. On the influence of the earth's rotation on ocean currents. *Ark. Mat. Astron. Fys.* 2 (11), 1–52.
- Fisher, J.C., Newton, R.J., Dila, D.K., McLellan, S.L., 2015. Urban microbial ecology of a freshwater estuary of Lake Michigan. *Elem. Sci. Anth.* 3 (64). <http://doi.org/10.12952/journal.elementa.000064>.
- Geyer, W.R., Ralston, D.K., 2015. Estuarine frontogenesis. *J. Phys. Oceanogr.* 45 (2), 546–561. <http://dx.doi.org/10.1175/JPO-D-14-0082.1>.
- Grunert, Brice, 2013. Evaluating the summer thermal structure of southern green bay, lake Michigan. Theses and Dissertations 252. <https://dc.uwm.edu/etd/252>.
- Hecky, R.E., Campbell, P., Hendzel, L.L., 1993. The stoichiometry of carbon, nitrogen, and phosphorus in particulate matter of lakes and oceans. *Limnol. Oceanogr.* 38. <http://dx.doi.org/10.4319/lo.1993.38.4.0709>.
- Hoffman, B.M., Lukoyanov, D., Yang, Z.-Y., Dean, D.R., Seefeldt, L.C., 2014. Mechanism of nitrogen fixation by nitrogenase: the next stage. *Chem. Rev.* 114 (8), 4041–4062. <http://doi.org/10.1021/cr400641x>.
- Kantha, L.H., Clayton, C.A., 1994. An improved mixed layer model for geophysical applications. *J. Geophys. Res.* 99 (25), 235–266.
- Knights, D., Parks, K.C., Sawyer, A.H., David, C.H., Browning, T.N., Danner, K.M., Wallace, C.D., 2017. Direct groundwater discharge and vulnerability to hidden nutrient loads along the Great Lakes coast of the United States. *J. Hydrol.* 554, 331–341.
- Kilroy, G., Coxon, C., 2005. Temporal variability of phosphorus fractions in Irish karst springs. *Environ. Geol.* 47, 421–430.
- Larson, J.H., Trebitz, A.S., Steinman, A.D., Wiley, M.J., Mazur, M.C., Pebbles, V., Braun, H.A., Seelbach, P.W., 2013. Great Lakes rivermouth ecosystems: scientific synthesis and management implications. *J. Great Lake Res.* 39, 513–524.
- Lawrence, G., Pieters, R., Zaremba, L., Tedford, T., Gu, L., Greco, S., Hamblin, P., 2004. Summer exchange between Hamilton Harbour and lake Ontario. *Deep-Sea Res. Part II Top. Stud. Oceanogr.* 51, 475–487.
- Levin, L.A., Ekau, W., Gooday, A.J., Jorissen, F., Middelburg, J.J., Naqvi, S.W.A., Neira, C., Rabalais, N.N., Zhang, J., 2009. Effects of natural and human-induced hypoxia on coastal benthos. *Biogeosciences* 6, 2063–2098. <https://doi.org/10.5194/bg-6-2063-2009>.
- Lewandowski, J., Meinikmann, K., Nutzmann, G., Rosenberry, D.O., 2015. Groundwater – the disregarded component in lake water and nutrient budgets. Part 2: effects of groundwater on nutrients. *Hydrol. Process.* 29, 2922–2955.
- Liu, Q., Rothstein, L.M., Luo, Y., Ullam, D.S., Codiga, D.L., 2016a. Dynamics of the periphery current in Rhode Island Sound. *Ocean Model.* 105, 13–24. <https://doi.org/10.1016/j.ocemod.2016.07.001>.
- Liu, Q., Rothstein, L.M., Luo, Y., 2016b. Dynamics of the Block island sound estuarine plume. *J. Phys. Oceanogr.* 46 (5), 1633–1656. <https://doi.org/10.1175/JPO-D-15-0099.1>.
- Liu, Q., Rothstein, L.M., Luo, Y., 2017. A periodic freshwater patch detachment process from the Block Island Sound estuarine plume. *J. Geophys. Res. Oceans* 122, 570–586. <http://dx.doi.org/10.1002/2015JC011546>.
- Liu, Q., Chai, F., Dugdale, R., Chao, Y., Xue, H., Rao, S., Wilkerson, F., Farrara, J., Zhang, H., Wang, Z., Zhang, Y.J., 2018. San Francisco bay nutrients and plankton dynamics as simulated by a coupled hydrodynamic-ecosystem model. *Contin. Shelf Res.* <https://doi.org/10.1016/j.csr.2018.03.008>.
- Marko, K.M., Rutherford, E.S., Eadie, B.J., Johengen, T.H., Lansing, M.B., 2013. Delivery of nutrients and seston from the Muskegon River watershed to near shore Lake Michigan. *J. Great Lake Res.* 39, 672–681.
- Mitchell, D.F., Wagner, K.J., Asbury, C., 1988. Direct measurement of groundwater flow and quality as a lake management tool. *Lake Reservoir Manag.* 4, 169–178.
- Mortimer, C.H., 1984. Measures and models in physical limnology. In: *hydrodynamics of Lakes*. In: Hutter, K. (Ed.), CISM Courses and Lectures Nr 286. Springer Verlag, Vienna-New York, pp. 287–322.
- Mortimer, C.H., 2004. Lake Michigan in Motion. Earth-spin, and Human Activities. The University of Wisconsin Press, pp. 173–196.
- Pitcher, G.C., Figueiras, F.G., Hickey, B.M., Moita, M.T., 2010. The physical oceanography of upwelling systems and the development of harmful algal blooms. *Prog. Oceanogr.* 85 (1–2), 5–32. <http://doi.org/10.1016/j.pcean.2010.02.002>.
- Plattner, S., Mason, D.M., Leshkevich, G.A., Schwab, D.J., Rutherford, E.S., 2006. Classifying and forecasting coastal upwellings in Lake Michigan using satellite derived temperature images and buoy data. *J. Great Lake Res.* 32 (1), 63–76. ISSN 0380-1330. [https://doi.org/10.3394/0380-1330\(2006\)32\[63:CAFCUJ\]2.0.CO;2](https://doi.org/10.3394/0380-1330(2006)32[63:CAFCUJ]2.0.CO;2).
- Rabalais, Nancy, N., Eugene Turner, R., Wiseman Jr, William J., 2002. Gulf of Mexico hypoxia, A.K.A. “the dead zone”. *Annu. Rev. Ecol. Systemat.* 33 (1), 235–263.
- Robinson, C., 2015. Review on groundwater as a source of nutrients to the Great Lakes and their tributaries. *J. Great Lake Res.* 41 (4), 941–950. <http://dx.doi.org/10.1016/j.jglr.2015.08.001>.
- Roegner, G.C., Needoba, J.A., Baptista, A.M., 2011. Coastal upwelling supplies oxygen-depleted water to the Columbia River estuary. *PLoS One* 6 (4), e18672. <https://doi.org/10.1371/journal.pone.0018672>.
- Roegner, G.C., Hickey, B.M., Newton, J.A., Shanks, A.L., Armstrong, D.A., 2002. Wind-induced plume and bloom intrusion into Willapa Bay, Washington. *Limnol. Oceanogr.* 47 (4), 1033–1042. <https://doi.org/10.4319/lo.2002.47.4.1033>.
- Rong, Z., Li, M., 2012. Tidal effects on the bulge region of Changjiang River plume. *Estuar. Coast Shelf Sci.* 97, 149–160. <http://dx.doi.org/10.1016/j.ecss.2011.11.035>.
- Safaie, A., Litchman, E., Phanikumar, M.S., 2017. Evaluating the role of groundwater in circulation and thermal structure within a deep inland lake. *Adv. Water Resour.* 108, 310–327. <http://dx.doi.org/10.1016/j.advwatres.2017.08.002>.
- Sanford, L.P., Sellner, K.G., Breitbart, D.L., 1990. Covariability of dissolved oxygen with physical processes in the summertime Chesapeake Bay. *J. Mar. Res.* 48, 567–590.
- Schwab, D.J., 1983. Numerical simulation of low-frequency current fluctuations in Lake Michigan. *J. Phys. Oceanogr.* 13, 2213–2224.
- Schwab, D.J., Bennett, J.R., 1987. Lagrangian comparison of objectively analyzed and dynamically modeled circulation patterns in Lake Erie. *J. Great Lake Res.* 13, 515–541.
- Schwab, D.J., Bedford, K.W., 1994. Initial implementation of the Great Lakes Forecasting System: a real-time system for predicting lake circulation and thermal structure. *Water Pollut. Res. J. Can.* 29 (2/3), 203–220.
- Schwab, D.J., 1977. Internal free oscillations in Lake Ontario. *Limnol. Oceanogr.* 22, 700–708.
- Simpson, J.H., Brown, J., Matthews, J., Allen, G., 1990. Tidal straining, density currents, and stirring in the control of estuarine stratification. *Estuaries* 13 (2), 125–132.
- Sinha, E., Michalak, A.M., Balaji, V., 2017. Eutrophication will increase during the 21<sup>st</sup> century as a result of precipitation changes. *Science* 357, 405–408.
- Steinman, A.D., Ogdahl, M., Rediske, R., Ruetz III, C.R., Biddanda, B.A., Nemeth, L., 2008. Current status and trends in Muskegon Lake, Michigan. *J. Great Lake Res.* 34, 169–188.
- Tang, Z., Engel, B.A., Pijanowski, B.C., Lim, K.J., 2005. Forecasting land use change and its environmental impact at a watershed scale. *J. Environ. Manag.* 76 (2), 35–45.
- Tranvik, L.J., Downing John, A., Cotner James, B., Loiselle Steven, A., Striegl Robert, G.,



- Ballatore Thomas, J., Dillon Peter, Finlay Kerri, Fortino Kenneth, Knoll Lesley, B., Kortelainen Pirkko, V., Kutser Tiit, Larsen Soren, Laurion Isabelle, Leech Dina, M., Leigh, McCallister S., McKnight Diane, M., Melack John, M., Overholt Erin, Porter Jason, A., Prairie Yves, Renwick William, H., Roland Fabio, Sherman Bradford, S., Schindler David, W., Sobek Sebastian, Tremblay Alain, Vanni Michael, J., Verschoor Antonie, M., von Wachenfeldt Eddie, Weyhenmeyer Gesa, A., 2009. Lakes and reservoirs as regulators of carbon cycling and climate. *Limnol. Oceanogr.* 54. <http://dx.doi.org/10.4319/lo.2009.54.6.part.2.2298>.
- Umlauf, L., Burchard, H., 2003. A generic length-scale equation for geophysical turbulence models. *J. Mar. Res.* 6, 235–265.
- Wang, G., Wang, D., Trenberth, K.E., Erfanian, A., Yu, M., Bosilovich, M.G., Parr, D.T., 2017. The peak structure and future changes of the relationships between extreme precipitation and temperature. *Nat. Clim. Change* 7, 268–274. <http://dx.doi.org/10.1038/nclimate3239>.
- Weinke, A.D., Kendall, S.T., Kroll, D.J., Strickler, E.A., Weinert, M.E., Holcomb, T.M., Defore, A.A., Dila, D.K., Snider, M.J., Gereaux, L.C., Biddanda, B.A., 2014. Systematically variable planktonic carbon metabolism along a land-to-lake gradient in a Great Lakes coastal zone. *J. Plankton Res.* 36, 1528–1542.
- Weinke, A.D., Biddanda, B.A., 2018. From bacteria to fish: ecological consequences of seasonal hypoxia in a Great Lakes estuary. *Ecosystems* 21 (3), 426–442. <https://doi.org/10.1007/s10021-017-0160-x>.
- Wiley, M.J., Hyndman, D.W., Pijanowski, B.C., et al., 2010. A multi-modeling approach to evaluating climate and land use change impacts in a Great Lakes River Basin. *Hydrobiologia* 657 (243). <https://doi.org/10.1007/s10750-010-0239-2>.
- Zhang, Y., Baptista, A.M., 2008. SELFE: a semi-implicit Eulerian-Lagrangian finite-element model for cross-scale ocean circulation. *Ocean Model.* 21 (3–4), 71–96.
- Zhang, Y.J., Witter, R.C., Priest, G.R., 2011. Tsunami-tide interaction in 1964 prince william sound tsunami. *Ocean Model.* 40 (3–4), 246–259. <https://doi.org/10.1016/j.ocemod.2011.09.005>.
- Zhang, Y., Ateljevich, E., Yu, H.-C., Wu, C.-H., Yu, J.C.S., 2015. A new vertical coordinate system for a three-dimensional unstructured-grid model. *Ocean Model.* 85, 16–31.
- Zhang, Y., Ye, F., Stanev, E.V., Grashorn, S., 2016. Seamless cross-scale modeling with SCHISM. *Ocean Model.* 102, 64–81. <http://dx.doi.org/10.1016/j.ocemod.2016.05.002>.
- Zhu, J., Weisberg, R.H., Zheng, L.Y., Han, S., 2014. Influences of channel deepening and widening on the tidal and non-tidal circulation of Tampa Bay. *Estuar. Coasts* 38, 132–150. <http://dx.doi.org/10.1007/s12237-014-9815-4>.